

## A Febrile Blood Donor

Julian W. Tang,<sup>1\*</sup> Yvonne Ng,<sup>2</sup> Evelyn S. C. Koay,<sup>1</sup> Gek Har Leow,<sup>1</sup> Eng Soo Yap,<sup>3</sup> Douglas Chan,<sup>1</sup>  
Lip Kun Tan,<sup>3</sup> and Paul A. Tambyah<sup>4</sup>

Departments of 1 Laboratory Medicine, 2 Neonatology, 3 Haematology and 4 Medicine,  
National University Hospital, Singapore.


\* Address correspondence to this author at: Department of Laboratory Medicine,  
National University Hospital, 5 Lower Kent Ridge Road, Singapore 119074,  
Singapore. Fax \_65-67771613; e-mail jvtang49@hotmail.com.

### CASE DESCRIPTION

A 4-day-old neonate born prematurely at 29 weeks gestation developed thrombocytopenia (platelet count  $43 \times 10^9/L$ ) with associated severe pulmonary and intracranial hemorrhaging. Urgent transfusions with 10 mL/kg of packed red cells and 10 mL/kg of platelet concentrate were given. These were sourced from blood donated 2 days earlier in Singapore by a 21-year old female university student who was clinically well at the time. The day after these blood products were given to the neonate, the blood donor contacted the blood transfusion service to inform them that she was now sick with a febrile flu-like illness. Her blood donation screening plasma sample was retrieved and tested by PCR for several viruses, including dengue, chikungunya, enterovirus, and Epstein-Barr virus.

Dengue RNA (serotype 2) was found to be present at a very low amount (i.e., at a real-time PCR cycle  $C_t$  number of 39–40) (Fig. 1A) from the initial RNA extract (extract A). To confirm the presence of this low amount of dengue 2 RNA, the same extract (extract A, which had been stored at  $-20^\circ C$ ) was retested (Fig. 1B), and again found to give a low-positive result.

| Questions to Consider  |
|--|
| <ul style="list-style-type: none"><li>• What steps should be taken next?</li></ul>                               |
| <ul style="list-style-type: none"><li>• What can cause false-negative and false-positive PCR results?</li></ul>  |
| <ul style="list-style-type: none"><li>• How should low-positive PCR results affect patient management?</li></ul> |



## Final Publication and Comments

The final published version with discussion and comments from the experts will appear in the March 2010 issue of *Clinical Chemistry*. To view the case and comments online, go to <http://www.clinchem.org/content/vol56/issue3> and follow the link to the Clinical Case Study and Commentaries.

## Educational Centers

If you are associated with an educational center and would like to receive the cases and questions 2-3 weeks in advance of publication, please email [clinchem@aacc.org](mailto:clinchem@aacc.org).

AACC is pleased to allow free reproduction and distribution of this Clinical Case Study for personal or classroom discussion use. When photocopying, please make sure the DOI and copyright notice appear on each copy.

All previous Clinical Cases Studies can be accessed and downloaded online at <http://www.aacc.org/resourcecenters/casestudies/>.

---

AACC is a leading professional society dedicated to improving healthcare through laboratory medicine. Its nearly 10,000 members are clinical laboratory professionals, physicians, research scientists, and others involved in developing tests and directing laboratory operations. AACC brings this community together with programs that advance knowledge, expertise, and innovation. AACC is best known for the respected scientific journal, *Clinical Chemistry*, the award-winning patient-centered web site *Lab Tests Online*, and the world's largest conference on laboratory medicine and technology. Through these and other programs, AACC advances laboratory medicine and the quality of patient care.