
Hemolytic Anemia Following Attempted Suicide

Nichole L. Korpi-Steiner¹, Jonathan B. Hoyne¹, James D. Hoyer¹, and Amy K. Saenger^{1*}

¹Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, MN

*Address correspondence to this author at: Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, 200 1st Street, SW, Rochester, MN 55905. Fax 507-538-7060; email saenger.amy@mayo.edu.

CASE DESCRIPTION

A 43-year-old African-American man with a history of hypertension, depression, and chronic alcohol abuse presented to the emergency service of an outside hospital complaining of chest pain. Laboratory testing indicated hepatocellular injury with aspartate aminotransferase (AST)² of 14000 U/L (reference interval 5–41 U/L) and alanine aminotransferase (ALT) of 6400 U/L (reference interval 8–45 U/L). Because laboratory indicators of liver function worsened, the patient was reinterviewed and confessed to having attempted suicide 3 days prior by ingesting about one-half gallon (approximately 4 L) of vodka and one-half bottle of extra-strength acetaminophen.

The patient was administered a bolus of N-acetylcysteine (NAC) at the outside hospital and referred to our institution because of concern for progression to fulminant hepatic failure. Laboratory tests were repeated and results were consistent with significant hepatocellular damage secondary to acetaminophen poisoning, including increased AST of 11000 U/L (reference interval 8–48 U/L), ALT of 6510 U/L (reference interval 7–55 U/L), total bilirubin of 395 μ mol/L (23.1 mg/dL) [reference interval 2–17 μ mol/L (0.1–1.0 mg/dL)], and direct-bilirubin of 207 μ mol/L (12.1 mg/dL) [reference interval <5.0 μ mol/L (<0.3 mg/dL)]. Treatment with intravenous infusion of NAC was continued, and nitroglycerin was administered for chest pain.

Within 24 h of admission the patient exhibited dyspnea with low hemoglobin (Hb) oxygen saturation (89%). Chest x-ray findings were unremarkable, whereas laboratory tests indicated a borderline-low Hb of 137 g/L (13.7 g/dL) [reference interval 135–175 g/L (13.5–17.5 g/dL)] with 4% methemoglobin (reference interval <1.5%). The patient received high-dose oxygen treatment but remained hypoxemic.

At 48 h the patient continued to have pronounced hyperbilirubinemia with evidence of hemolytic anemia, which included low Hb of 7.5 g/dL with 6% methemoglobin, marked reticulocytosis of 11.5% (reference interval 0.6%–1.83%), nondetectable haptoglobin [1.4 μ mol/L (<14 mg/dL); reference interval 3–20 μ mol/L (30–200 mg/dL)], increased mean corpuscular volume (MCV) (97.8 fL; reference interval 81.2–95.1 fL), and increased lactate dehydrogenase (LD) (6870 U/L; reference interval 122–222 U/L). Moreover, results of a direct Coombs test were negative, although a routine peripheral blood smear revealed the presence of “bite cells.” An enzymopathy with a drug-induced acute episode of hemolytic anemia was suspected.

Questions to Consider
<ul style="list-style-type: none">• How does acetaminophen poisoning disrupt typical acetaminophen metabolism and what are the consequences?
<ul style="list-style-type: none">• What are potential causes of hemolytic anemia?
<ul style="list-style-type: none">• What laboratory tests should be performed to diagnose hemolytic anemia?

Final Publication and Comments

The final published version with discussion and comments from the experts will appear in the December 2008 issue of *Clinical Chemistry*. To view the case and comments online, go to <http://www.clinchem.org/content/vol54/issue12/> and follow the link to the Clinical Case Study and Commentaries.

Educational Centers

If you are associated with an educational center and would like to receive the cases and questions 2-3 weeks in advance of publication, please email clinchem@aacc.org.

AACC is pleased to allow free reproduction and distribution of this Clinical Case Study for personal or classroom discussion use. When photocopying, please make sure the DOI and copyright notice appear on each copy.

All previous Clinical Cases Studies can be accessed and downloaded online at <http://www.aacc.org/resourcecenters/casestudies/>.

AACC is a leading professional society dedicated to improving healthcare through laboratory medicine. Its nearly 10,000 members are clinical laboratory professionals, physicians, research scientists, and others involved in developing tests and directing laboratory operations. AACC brings this community together with programs that advance knowledge, expertise, and innovation. AACC is best known for the respected scientific journal, *Clinical Chemistry*, the award-winning patient-centered web site *Lab Tests Online*, and the world's largest conference on laboratory medicine and technology. Through these and other programs, AACC advances laboratory medicine and the quality of patient care.