

Scenario 1	Δ Triglycerides	Total Cholesterol (mg/dL)	HDL-C (mg/dL)	Triglycerides (mg/dL)	LDL-C* (mg/dL)	Non-Fasting LDL-C Change		Framingham 10-yr Risk**	Pooled Cohort**
						(mg/dL)	%		
0% (Fasting)	240	39	195	162	-	-	9.0%	3.9%	
+10%			215	158	-4	-2.4%			
+20%			234	154	-8	-4.8%			
+50%			293	143	-19	-12.0%			
Scenario 2	Δ Triglycerides	Total Cholesterol (mg/dL)	HDL-C (mg/dL)	Triglycerides (mg/dL)	LDL-C* (mg/dL)	Non-Fasting LDL-C Change		Framingham 10-yr Risk**	Pooled Cohort**
						(mg/dL)	%		
0% (Fasting)	250	30	132	194	-	-	11.4%	5.0%	
+10%			145	191	-3	-1.3%			
+20%			158	188	-5	-2.7%			
+50%			198	180	-13	-6.7%			

*LDL-C estimated by the Friedewald formula

**Risk scores calculated using a hypothetical 58-year-old female non-smoker with systolic blood pressure of 125 mmHg and C-reactive protein of 1.0 mg/L.