

---

## Unusual Cause of Abdominal Pain and Anemia

Cilie C. van 't Klooster,<sup>1\*</sup> Jan J. Uil,<sup>1</sup> Joep van der Leeuw,<sup>1</sup> Elaine F. Eppens,<sup>1</sup> and Susanne C. Marczinski<sup>1</sup>

<sup>1</sup>Ziekenhuis Gelderse Vallei, Ede, the Netherlands.

\*Address correspondence to this author at: UMC Utrecht, Heidelberglaan 100, 3584 CX, Utrecht, the Netherlands. Fax 887-555-639; e-mail [cilie.vantklooster@gmail.com](mailto:cilie.vantklooster@gmail.com)

---

### CASE DESCRIPTION

A 46-year-old man, originally from Iran, with no relevant medical history, was admitted with intermittent acute colic pain in the lower abdomen with nausea and vomiting over the past few weeks. On physical examination, the patient had lower abdominal pain without guarding. Laboratory findings showed a microcytic anemia, a heterozygote  $\beta$ -thalassemia, mild leukocytosis, and slight liver enzyme activity increase (Table 1, case 1). Hepatitis, cytomegalovirus, and Epstein-Barr virus were negative. Ultrasonography and computed tomography scan of the abdomen did not suggest liver, kidney, or pancreas disease, and gastroscopy and colonoscopy revealed no pathology. The patient denied substance abuse. The symptoms improved spontaneously and the patient awaited further diagnostics in the outpatient clinic. Within 1 week, the patient returned with similar symptoms. Acute intermittent porphyria (AIP)<sup>2</sup> was suspected and urine porphyrins were measured. Urine  $\delta$ -aminolevulinic acid (ALA) concentration was increased at 96.9 mmol/mol of creatinine (reference interval <3.9 mmol/mol creatinine), porphobilinogen (PBG) concentration was normal at 8  $\mu$ mol/L (reference interval 0–9  $\mu$ mol/L), and coproporphyrinogen III concentration was increased at 91.8 nmol/mmol of creatinine (reference interval 2.9–19.3 nmol/mmol of creatinine). These findings excluded the diagnosis AIP, as the PBG concentrations remained normal and both ALA concentration and coproporphyrinogen III concentrations increased.

Another 42-year-old man, originally from Iran and with no relevant medical history, presented with similar complaints of acute diffuse colic abdominal pain with nausea and vomiting. He complained of diffuse abdominal pain without guarding. Laboratory findings showed microcytic anemia, without signs of iron deficiency or thalassemia; basophilic stippling of red blood cells; mild leukocytosis; and increased liver enzyme activities (Table 1, case 2). Viral serology was negative. The patient denied substance abuse. Considering a gastric ulcer, a gastroscopy was performed that showed no abnormalities. Doppler ultrasonography of the abdomen was negative for liver, kidney, or pancreas disease, or mesenteric ischemia. On the basis of the symptoms, porphyrin analysis was performed, with the following results: ALA concentration was increased at 79.6 mmol/mol of creatinine and the free erythrocyte protoporphyrin concentration was also increased at 126.3  $\mu$ g/dL (reference interval 0–65  $\mu$ g/dL). The patient's gums revealed Burton's lines. Unknown to the medical team at the time, the 2 patients were friends.

Soon after, a third patient, sibling of the aforementioned second case, presented with similar complaints, and the laboratory tests revealed similar results (Table 1, case 3).

QUESTIONS TO CONSIDER	
•	What are the main causes of microcytic anemia?
•	What are the diagnostic tests for acute intermittent porphyria?
•	Which secondary porphyria can mimic acute porphyria?

	Reference interval	Case 1	Case 2	Case 3
Hemoglobin (g/dl)	Men 14.0-17.5 Women 12.3-15.3	10.3	8.4	9.5
Mean corpuscular volume (fL)	80-100	60	78	78
White blood cell count (/nL)	4.0-11.0	11.3	12.5	5.0
Red blood cell	–	Basophilic stippling	Basophilic stippling	Basophilic stippling
Gamma glutamyl transpeptidase (IU/L)	0-55	57	354	49
Alkaline phosphatase (IU/L)	0-120	121	197	91
Alanine aminotransferase (IU/L)	0-45	71	91	27

### Final Publication and Comments

The final published version with discussion and comments from the experts will appear in the December 2017 issue of *Clinical Chemistry*. To view the case and comments online, go to <http://www.clinchem.org/content/vol63/issue12> and follow the link to the Clinical Case Study and Commentaries.

### Educational Centers

If you are associated with an educational center and would like to receive the cases and questions 1 month in advance of publication, please email [clinchemed@aacc.org](mailto:clinchemed@aacc.org).

All previous Clinical Case Studies can be accessed and downloaded online at <https://www.aacc.org/publications/clinical-chemistry/clinical-case-studies>

AACC is pleased to allow free reproduction and distribution of this Clinical Case Study for personal or classroom discussion use. When photocopying, please make sure the DOI and copyright notice appear on each copy.

---

AACC is a leading professional society dedicated to improving healthcare through laboratory medicine. Its nearly 10,000 members are clinical laboratory professionals, physicians, research scientists, and others involved in developing tests and directing laboratory operations. AACC brings this community together with programs that advance knowledge, expertise, and innovation. AACC is best known for the respected scientific journal, *Clinical Chemistry*, the award-winning patient-centered web site *Lab Tests Online*, and the world's largest conference on laboratory medicine and technology. Through these and other programs, AACC advances laboratory medicine and the quality of patient care.